

INSTRUCTIONS FOR 2 DAY COLONOSCOPY PREP

MiraLAX with Gatorade/PowerAde/Magnesium Citrate

***You will need to purchase all of the following over the counter items:

MiraLAX 17.9oz (510 gram) bottle over the counter, a Total of 64oz of Gatorade or PowerAde (no red or purple), Dulcolax laxative tablets 5mg (total of 40mg) and 2 bottles of Magnesium Citrate***

Colonoscopy: The visual examination of the large intestine (colon) using a lighted flexible fiber optic endoscope.

Note: Individual responses to laxatives do vary. This prep may cause multiple bowel movements. Often works within 30 minutes but may take as long as 3 hours. If you are unable to keep the prep down, please call your physician.

Consult your Primary Care Physician if you are taking or have the following medical concerns:

Coumadin/Jantoven/warfarin: Stop it five (5) days before your colonoscopy if approved by your physician prescribing this medication.

Do not take this medication on the day of your procedure. Ask your physician when to resume this medication

Plavix (clopidogrel)/Pradaxa/Pletal (cilostazol)/Effient and Eliquis; Stop it three (3) days before colonoscopy-**only if approved by prescribing physician**

Brilinta: Stop five (5) days before colonoscopy- **only if approved by prescribing physician**

Aggrenox: Stop three (3) days to five (5) days prior to colonoscopy- **only if approved by prescribing physician**

Xarelto/Savaysa: Stop three (3) days before colonoscopy- **only if approved by prescribing physician**

Iron pills: Stop for seven (7) days before colonoscopy

Dialysis Treatment: Alternate treatment may be used.

****Anti-Seizure Medication:** Take at least 2 hours before the scheduled colonoscopy with a small sip of water.

****Blood Pressure/Heart Medication:** Please take your morning dose at least 2 hours before your colonoscopy with a small sip of water.

Insulin: You may need to adjust your insulin dosages. Please speak with your monitoring physician for further instructions. **Ask your physician when to resume your insulin.**

Inhalers: Please bring your "rescue" inhaler to your appointment, even if you have not needed to use it for some time.

Heart Valve Replacement/ Pacemaker with Internal Defibrillator

Semaglutide medication will need to be stopped 7 days prior to procedure.

*Dulaglutide (Trulicity) *Exenatide (Byetta) *Exenatide ER *Bydureon BCise *Liraglutide (Victoza) *Lixisenatide, *Semaglutide subcutaneous/tablet *Ozempic *Rybelsus *Tirzepatide (Mounjaro) *Wegovy

TWO DAYS BEFORE COLONOSCOPY

MAY HAVE A LIGHT BREAKFAST, LIQUIDS THE REMAINDER OF THE DAY

No tomatoes or corn 2 days prior to procedure

At 4 PM TAKE 4 (20mg total) Dulcolax Laxative Tablets with a full glass of water

At 6 PM Drink 1 bottle of Magnesium Citrate with at least 16 oz of clear liquids

ONE DAY BEFORE COLONOSCOPY

If you have been holding your Coumadin/Jantoven/Warfarin you will need a PT/INR (lab test) taken 24 hours prior to exam if requested by physician. **CLEAR LIQUIDS DIET ALL DAY – no milk or milk products**

At 4 PM TAKE 4 (20mg total) Dulcolax Laxative Tablets and 1 bottle of Magnesium Citrate with at least 16 oz of clear liquids

**** Mix 64oz of Gatorade with 14 capfuls of MiraLAX**

At 6 PM Drink half of the MiraLAX/Gatorade mixture. (You do not have to drink it all at once, but finish by 9pm for next dose)

At 9 PM Drink the rest of the MiraLAX/Gatorade mixture. (Finish before midnight)

CLEAR LIQUIDS

Water	Strained fruit juices without pulp	Broth or Bouillon
Jello – any flavor – no red jello	Clear sodas – 7up, Sprite, Ginger Ale, etc.	Popsicles – no red or purple
Coffee or Tea without creamer	Ensure and Ensure plus (vanilla only)	No red or purple liquids

DAY OF COLONOSCOPY

1. Nothing by mouth after midnight except approved **medications listed above**** with a sip of water. You may brush your teeth and rinse with water. **NO GUM, MINTS OR LOZENGES. NO CHEWING TOBACCO**

2. **You will be medicated and you need to bring someone with you to drive you home.**

3. Bring a current list of medications and colostomy supplies if applicable; if Menstruating please use a tampon if possible.

On the day of your procedure, report to the following unless otherwise instructed by your physician:

The Endoscopy Center of Topeka, 785.354.1254

2200 SW 6th Avenue, Suite 103

Topeka, KS 66606

Check-in Date: _____

Check-in Time: _____

MKC Gastroenterology, 785-354-8518

If you are unable to keep your appointment, please notify the office within 48 hours prior to your scheduled procedure.